

Epilobium parviflorum

Habitus and growth type

Plant height [m]: **0.48**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **29.78**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **8.5**

Reaction indicator value: **7**

Nutrient indicator value: **5.5**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.01**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.48**

Grazing pressure: **0.25**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[QG \(EPI\) *Epilobietea angustifolii*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline),
Saline vegetation, Synanthropic**

Distribution

Continentality amplitude: **5**