

Erica manipuliflora

Habitus and growth type

Plant height [m]: **0.45**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Spinescence: **not spinescent**

Flower

Flowering period: **July-September**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.48**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.02**

Grazing pressure: **0.31**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#)

Constant species of EUNIS habitats: [S63 Eastern garrigue](#)

Dominant species of EUNIS habitats: [S63 Eastern garrigue](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**