Erigeron atticus

Habitus and growth type

Plant height [m]: **0.36**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.24**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **2.5**
Moisture indicator value: **5**
Reaction indicator value: **6**
Nutrient indicator value: **3.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.62**
Disturbance frequency (herb layer): **0.85**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.28**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.14**
Grazing pressure: **0.16**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea***

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Forest**

Distribution

Continentality: **5**
Continentality amplitude: **1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz