

Erodium foetidum

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **5.1**

Moisture indicator value: **3.2**

Reaction indicator value: **6.2**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **1.15**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.13**

Grazing pressure: **0.31**

Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1H Iberian oromediterranean basiphilous dry grassland](#), [S73 Western Mediterranean mountain hedgehog-heath](#), [U34 Mediterranean siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**