

Erodium laciniatum

Habitus and growth type

Plant height [m]: **0.12**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.07**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **10**

Moisture indicator value: **2.5**

Reaction indicator value: **7.5**

Nutrient indicator value: **7**

Salinity indicator value: **3**

Disturbance indicator values

Disturbance frequency: **1.26**

Disturbance frequency (herb layer): **1.73**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.1**
Grazing pressure: **0.26**
Soil disturbance: **0.43**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FH \(BUL\) *Poetea bulbosae*](#), [FI \(TUB\) *Helianthemetea guttati*](#), [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N16 Mediterranean and Macaronesian coastal dune grassland \(grey dune\)](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**