

Eruca vesicaria

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **16.9**

Flower

Flowering period: **January-June**

Fruit, seed and dispersal

Seed mass [mg]: **1.43**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**

Moisture indicator value: **4.2**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **2.03**

Disturbance severity: **0.8**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.72**

Grazing pressure: **0.21**

Soil disturbance: **0.51**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality amplitude: **5**