Erythronium dens-canis

Habitus and growth type

Plant height [m]: **0.18**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **1.62**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **3.8**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Nutrient indicator value: **3.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.07**
Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.02**
Grazing pressure: **0.21**
Soil disturbance: **0.1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz