

# ***Euphorbia platyphyllos***

## **Habitus and growth type**

Plant height [m]: **0.42**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **May-September**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.21**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **6.8**

Moisture indicator value: **5.1**

Reaction indicator value: **6.5**

Nutrient indicator value: **5.8**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **0.79**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.62**

Mowing frequency: **0.4**

Grazing pressure: **0.22**

Soil disturbance: **0.46**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N1E Black Sea broad-leaved coastal dune forest](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Wetland, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality amplitude: **5**