

Euphorbia prostrata

Habitus and growth type

Plant height [m]: **0.08**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **May-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.15**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **9**

Moisture indicator value: **2.5**

Reaction indicator value: **6**

Nutrient indicator value: **4.5**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.82**

Disturbance frequency (herb layer): **1.82**

Disturbance severity: **0.85**

Disturbance severity (herb layer): **0.85**

Mowing frequency: **0.96**

Grazing pressure: **0.11**

Soil disturbance: **0.58**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) *Digitario sanguinalis*-*Eragrostietea minoris*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V34 Trampled xeric grassland with annuals](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**