

Fallopia dumetorum

Habitus and growth type

Plant height [m]: **1.34**

Life span: **Annual**

Life form: **Therophyte, Herbaceous liana**

Leaf

Specific leaf area [mm²/mg]: **34.58**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **3.74**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**

Moisture indicator value: **5.1**

Reaction indicator value: **6.2**

Nutrient indicator value: **6.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.22**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.11**

Grazing pressure: **0.19**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) *Epilobietea angustifolii*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**