Fallopia dumetorum

Habitus and growth type

Plant height [m]: **1.34**
Life span: **Annual**
Life form: **Therophyte, Herbaceous liana**

Leaf

Specific leaf area [mm^2/mg]: **34.58**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **3.74**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**
Moisture indicator value: **5.1**
Reaction indicator value: **6.2**
Nutrient indicator value: **6.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.22**
Disturbance frequency (herb layer): **1.34**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.11**
Grazing pressure: **0.19**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz