Ferulago campestris

Habitus and growth type

Plant height [m]: **0.8**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Moisture indicator value: **4.3**
Reaction indicator value: **7.5**
Nutrient indicator value: **6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.55**
Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.35**
Grazing pressure: **0.29**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz