

# ***Festuca braun-blanquetii***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **8**

Moisture indicator value: **6**

Reaction indicator value: **4**

Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CN \(SAB\) Junipero-Pinetea sylvestris](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**