

Festuca richardsonii

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Moisture indicator value: **5**

Disturbance indicator values

Disturbance frequency: **1.61**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.26**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.02**

Grazing pressure: **0.23**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**AE \(COC\) Saxifrago cernuae-Cochlearieteae groenlandicae**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**MA211 Arctic coastal saltmarsh**](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Alpine-subalpine and arctic grassland**