

Fourraea alpina

Habitus and growth type

Plant height [m]: **0.61**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.3**

Temperature indicator value: **5.6**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.16**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.18**

Mowing frequency: **0.06**

Grazing pressure: **0.21**

Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[CB \(PUB\) *Quercetea pubescentis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**

Continentality amplitude: **2**