

Fumana ericoides

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **5.22**

Fruit, seed and dispersal

Seed mass [mg]: **2.44**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **9**

Disturbance indicator values

Disturbance frequency: **0.65**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.11**

Grazing pressure: **0.38**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S61 Western basiphilous garrigue](#), [S65 Mediterranean gypsum scrub](#)

Constant species of EUNIS habitats: [S61 Western basiphilous garrigue](#), [S65 Mediterranean gypsum scrub](#)

Broad habitat

Occurrence in broad habitats: **Scrub**