

# ***Fumaria capreolata***

## **Habitus and growth type**

Plant height [m]: **0.63**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **January-May, November-December**

## **Fruit, seed and dispersal**

Seed mass [mg]: **3.58**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7**

Moisture indicator value: **4**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.79**

Disturbance severity (herb layer): **0.66**

Mowing frequency: **0.45**

Grazing pressure: **0.23**

Soil disturbance: **0.41**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1L Madeiran oromediterranean siliceous dry grassland](#)

Constant species of EUNIS habitats: [R1L Madeiran oromediterranean siliceous dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

## Distribution

Continentality amplitude: **5**