

# *Galium parisiense*

## Habitus and growth type

Plant height [m]: **0.21**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.47**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.19**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7.5**

Moisture indicator value: **2.3**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.04**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.37**

Grazing pressure: **0.33**

Soil disturbance: **0.31**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#), [QA \(PAR\) \*Papaveretea rhoeadis\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

Constant species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality amplitude: **5**