Galium parisiense

Habitus and growth type

Plant height [m]: **0.21**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **24.47**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.19**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **7.5**
Moisture indicator value: **2.3**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.04**
Disturbance frequency (herb layer): **1.85**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.37**
Grazing pressure: **0.33**
Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati*, QA (PAR) *Papaveretea rhoeadis*, QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**
Constant species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz