Genista florida

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Seed mass [mg]: **7.45**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **4**
Reaction indicator value: **2**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.21**
Mowing frequency: **0**
Grazing pressure: **0.28**
Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S33 Lowland to montane temperate and submediterranean genistoid scrub**
Constant species of EUNIS habitats: **S33 Lowland to montane temperate and**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

submediterranean genistoid scrub

Dominant species of EUNIS habitats: **S33 Lowland to montane temperate and submediterranean genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz