

Gentiana occidentalis

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.35**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **3**

Moisture indicator value: **4.5**

Reaction indicator value: **7**

Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.58**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.18**

Grazing pressure: **0.3**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) *Elyno-Seslerietea*](#), [FP \(ONO\) *Festuco hystricis-Ononidetea striatae*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**