Gentianella anisodonta

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Flower

Flowering period: **May-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **2.7**
Moisture indicator value: **4**
Reaction indicator value: **7**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.72**
Disturbance frequency (herb layer): **0.91**
Disturbance severity: **0.33**
Disturbance severity (herb layer): **0.26**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.04**
Grazing pressure: **0.22**
Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz