

Geranium sanguineum

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **21.83**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **9.46**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **6**

Moisture indicator value: **3.3**

Reaction indicator value: **7.4**

Nutrient indicator value: **2.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**

Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.23**
Grazing pressure: **0.24**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#)

Constant species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#), [S36 Low steppic scrub](#), [T36 Temperate and submediterranean montane *Pinus sylvestris*-*Pinus nigra* forest](#)

Dominant species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **5**