Geum coccineum

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.3**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Temperature indicator value: **6**
Moisture indicator value: **7**

Disturbance indicator values

Disturbance frequency: **0.92**
Disturbance frequency (herb layer): **1.02**
Disturbance severity: **0.32**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.24**
Grazing pressure: **0.15**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CS (MUL) *Mulgedio-Aconitetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q23 Relict mire of Mediterranean mountains**
Constant species of EUNIS habitats: **Q23 Relict mire of Mediterranean mountains**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Mire, Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz