# *Glandora prostrata*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6.5**
Moisture indicator value: **4**
Reaction indicator value: **2.5**
Nutrient indicator value: **1.5**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.43**
Disturbance frequency (herb layer): **1.35**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.07**
Grazing pressure: **0.28**
Soil disturbance: **0.12**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**, **T29 Broadleaved evergreen plantation of non site-native trees**

### Broad habitat

Occurrence in broad habitats: **Heathland, Scrub, Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*