

Globularia bisnagarica

Habitus and growth type

Plant height [m]: **0.14**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **8.22**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.66**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **6.3**

Moisture indicator value: **2.4**

Reaction indicator value: **8.6**

Nutrient indicator value: **1.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.99**

Disturbance frequency (herb layer): **1.78**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0.42**

Grazing pressure: **0.33**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**DA \(FES\) Festuco-Brometea, FP
\(ONO\) Festuco hystricis-Ononidetea striatae**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**](#)

Constant species of EUNIS habitats: [**R16 Perennial rocky grassland of Central and South-Eastern Europe, R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **4**