

# *Glyceria maxima*

## Habitus and growth type

Plant height [m]: **1.65**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **19.79**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.76**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **5.3**

Moisture indicator value: **9.8**

Reaction indicator value: **7.7**

Nutrient indicator value: **8.1**

Salinity indicator value: **0.1**

### Disturbance indicator values

Disturbance frequency: **1.42**

Disturbance frequency (herb layer): **1.73**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.38**

Grazing pressure: **0.15**

Soil disturbance: **0.1**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [\*\*OD \(PHR\) \*Phragmito-Magnocaricetea\*\*\*](#)

### **EUNIS habitat**

Diagnostic species of EUNIS habitats: [\*\*Q51 Tall-helophyte bed\*\*](#)

Constant species of EUNIS habitats: [\*\*P3b Fresh-water small pleustophyte vegetation, Q51 Tall-helophyte bed, Q52 Small-helophyte bed, Q53 Tall-sedge bed, T15 Broadleaved swamp forest on non-acid peat\*\*](#)

### **Broad habitat**

Occurrence in broad habitats: **Aquatic, Wetland, Forest**

## **Distribution**

Continentality amplitude: 7