

Glycyrrhiza glabra

Habitus and growth type

Plant height [m]: **0.84**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **6.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Reaction indicator value: **7.2**

Nutrient indicator value: **6.4**

Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.3**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.56**

Mowing frequency: **0.41**

Grazing pressure: **0.2**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#), [DB \(FEP\) *Festuco-Puccinellietea*](#), [EB \(TAM\) *Tamaricetea arceuthoidis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N35 Mediterranean and Black Sea soft sea cliff](#), [R65 Continental subsaline alluvial pasture and meadow](#), [S68 Semi-desert sand dune with sparse scrub](#)

Constant species of EUNIS habitats: [N35 Mediterranean and Black Sea soft sea cliff](#), [R65 Continental subsaline alluvial pasture and meadow](#), [S68 Semi-desert sand dune with sparse scrub](#)

Dominant species of EUNIS habitats: [N35 Mediterranean and Black Sea soft sea cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff, Grassland (non-alpine, non-saline), Saline vegetation, Semi-desert, Scrub, Synanthropic**