

# *Gnaphalium norvegicum*

## Habitus and growth type

Plant height [m]: **0.22**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **21.17**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.09**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **2.8**

Moisture indicator value: **4.7**

Reaction indicator value: **4.2**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.83**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.06**

Grazing pressure: **0.24**

Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CS \(MUL\) \*Mulgedio-Aconitetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T1C Temperate and boreal mountain \*Betula\* and \*Populus tremula\* forest on mineral soils](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

## Distribution

Continentality: **5**

Continentality amplitude: **4**