Halimione verrucifera

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.8**
Moisture indicator value: **6.8**

Disturbance indicator values

Disturbance frequency: **1.91**
Disturbance frequency (herb layer): **2.09**
Disturbance severity: **0.45**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0.06**
Grazing pressure: **0.45**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **EC (KAL) *Kalidietea foliati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **MA241 Black Sea littoral saltmarsh**, **R64 Semi-desert salt pan**
Constant species of EUNIS habitats: **MA241 Black Sea littoral saltmarsh**, **R64 Semi-**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

desert salt pan

Dominant species of EUNIS habitats: **R64 Semi-desert salt pan**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz