Hedysarum boveanum

Habitus and growth type

Plant height [m]: **0.32**
Life span: **Perennial**
Life form: **Hemicryptophyte**
Spinescence: **not spinescent**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **5.9**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **7**
Moisture indicator value: **2.5**
Reaction indicator value: **7.5**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.65**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.19**
Mowing frequency: **0**
Grazing pressure: **0.44**
Soil disturbance: **0.19**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz