Helictochloa marginata

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6**
Moisture indicator value: **4**
Reaction indicator value: **3**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**
Disturbance frequency (herb layer): **1.5**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.18**
Grazing pressure: **0.31**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of EUNIS habitats: **T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest**

Constant species of EUNIS habitats: **T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz