Helichrysum litoreum

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Chamaephyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **9.6**
Moisture indicator value: **2.9**
Reaction indicator value: **6.8**
Nutrient indicator value: **3.5**
Salinity indicator value: **4**

Disturbance indicator values

Disturbance frequency: **0.61**
Disturbance frequency (herb layer): **0.7**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.09**
Grazing pressure: **0.13**
Soil disturbance: **0.06**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N32 Mediterranean and Black Sea rocky sea cliff and shore**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz