Helichrysum orientale

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **11**
Moisture indicator value: **2**
Reaction indicator value: **8**
Nutrient indicator value: **3**
Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.37**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.09**
Grazing pressure: **0.24**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KC (ASP) *Asplenietea trichomanis***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz