

Heliosperma pusillum

Habitus and growth type

Plant height [m]: **0.11**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.13**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **3.8**

Moisture indicator value: **8**

Reaction indicator value: **8.3**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.55**

Disturbance frequency (herb layer): **0.86**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.03**

Grazing pressure: **0.17**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) *Asplenietea trichomanis*](#),
[OA \(MON\) *Montio-Cardaminetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [P2N Spring](#)

Constant species of EUNIS habitats: [P2N Spring](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Spring, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **2**