Hierochloe odorata

Habitus and growth type

Plant height [m]: **0.46**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **41.51**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.9**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.2**
Temperature indicator value: **4.8**
Moisture indicator value: **7.7**
Reaction indicator value: **4.8**
Nutrient indicator value: **2.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.79**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.44**
Grazing pressure: **0.19**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Grassland (non-alpine, non-saline)**

Distribution

Continentality: **7**
Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz