Hippuris vulgaris

Habitus and growth type

Plant height [m]: **0.58**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **32.53**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.9**
Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Temperature indicator value: **5.1**
Moisture indicator value: **10.5**
Reaction indicator value: **7.3**
Nutrient indicator value: **4.8**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.7**
Disturbance frequency (herb layer): **1.71**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.27**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.1**
Grazing pressure: **0.1**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz