

# *Hylotelephium vulgare*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **5**

Moisture indicator value: **6.8**

Disturbance indicator values

Disturbance frequency: **0.15**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.06**

Grazing pressure: **0.15**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **4**