Hymenocarpos circinnatus

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **6.3**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **10**
Moisture indicator value: **2.5**
Reaction indicator value: **6.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.78**
Disturbance frequency (herb layer): **1.8**
Disturbance severity: **0.78**
Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.27**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.34**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**
Constant species of EUNIS habitats: **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz