

# *Hymenocarpus circinnatus*

## Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **February-April**

## Fruit, seed and dispersal

Seed mass [mg]: **6.3**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **2.5**

Reaction indicator value: **6.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.78**

Disturbance frequency (herb layer): **1.8**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.27**

Grazing pressure: **0.32**

Soil disturbance: **0.34**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*S72 Eastern Mediterranean spiny heath \(phrygana\)\*\*](#)

Constant species of EUNIS habitats: [\*\*S72 Eastern Mediterranean spiny heath \(phrygana\)\*\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**