

# *Hyoseris scabra*

## Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **January-April**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **4**

Reaction indicator value: **7.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.24**

Disturbance frequency (herb layer): **1.8**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.73**

Mowing frequency: **0.37**

Grazing pressure: **0.33**

Soil disturbance: **0.39**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) Stipo-Trachynietea distachyae](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**