Hypericum androsaemum

Habitus and growth type

Plant height [m]: **0.55**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.5**
Temperature indicator value: **7**
Moisture indicator value: **6.8**
Reaction indicator value: **5.5**
Nutrient indicator value: **5.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.14**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.19**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.02**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***, **CC (QUE) *Quercetea robori-petraeae***, **HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T22 Mainland laurophyllous forest**
Constant species of EUNIS habitats: **T22 Mainland laurophyllous forest**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz