

Hypericum linarifolium

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **6.5**

Moisture indicator value: **3**

Reaction indicator value: **3**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.89**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.19**

Grazing pressure: **0.34**

Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S62 Western acidophilous garrigue](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**