Chaerophyllum aureum

Habitus and growth type

Plant height [m]: **0.9**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **21.52**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **7.59**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Hypertrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.4**
Reaction indicator value: **8**
Nutrient indicator value: **8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.53**
Disturbance frequency (herb layer): **1.42**
Disturbance severity: **0.59**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.41**
Mowing frequency: **0.39**
Grazing pressure: **0.18**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CR (VIR) *Betulo carpaticae-Alnetea viridis*, QG (EPI) *Epilobietea angustifolii***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V39 Mesic perennial anthropogenic herbaceous vegetation**
Dominant species of EUNIS habitats: **V39 Mesic perennial anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz