

Chamaemelum nobile

Habitus and growth type

Plant height [m]: **0.24**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.19**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Reaction indicator value: **4**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.97**

Disturbance frequency (herb layer): **2.29**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.66**

Grazing pressure: **0.59**

Soil disturbance: **0.33**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#)

Constant species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**