

Chenopodiastrium hybridum

Habitus and growth type

Plant height [m]: **0.62**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **35.61**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **1.41**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **5.9**

Moisture indicator value: **5.1**

Reaction indicator value: **7.7**

Nutrient indicator value: **7.8**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **1.96**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.7**

Grazing pressure: **0.14**

Soil disturbance: **0.75**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) Sisymbrietea](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: **7**

Continentality amplitude: **8**