

Cherleria biflora

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **1**

Moisture indicator value: **4.3**

Reaction indicator value: **3**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.08**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0**

Grazing pressure: **0.15**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U21 Boreal and Arctic siliceous scree and block field](#)

Constant species of EUNIS habitats: [U21 Boreal and Arctic siliceous scree and block field](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**