

# *Cherleria circassica*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **11.89**

## Fruit, seed and dispersal

Seed mass [mg]: **0.46**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Disturbance indicator values

Disturbance frequency: **1.65**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0**

Grazing pressure: **0.25**

Soil disturbance: **0.07**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CT \(TRI\) \*Juncetea trifidi\*](#), [LA \(HER\) \*Salicetea herbaceae\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**