Chorispora tenella

Habitus and growth type

Plant height [m]: **0.24**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6.5**
Moisture indicator value: **3.8**

Disturbance indicator values

Disturbance frequency: **2.17**
Disturbance frequency (herb layer): **2.13**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.25**
Grazing pressure: **0.3**
Soil disturbance: **0.49**

Habitat and sociology

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**, **S68 Semi-desert sand dune with sparse scrub**

Constant species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**, **S68 Semi-desert sand dune with sparse scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz