Impatiens parviflora

Habitus and growth type

Plant height [m]: **0.51**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **144.78**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **8.1**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4**
Temperature indicator value: **5.8**
Moisture indicator value: **5.4**
Reaction indicator value: **6.6**
Nutrient indicator value: **6.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.16**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.07**
Grazing pressure: **0.19**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae*, QG (EPI) *Epilobietea angustifolii***

EUNIS habitat

Constant species of EUNIS habitats: **R57 Herbaceous forest clearing vegetation, T13 Temperate hardwood riparian forest**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz